

# Effects of short term exposure with mine tailings on deep-sea benthic ecosystem function and services

Mevenkamp Lisa<sup>1</sup>, Tanja Stratmann<sup>2</sup>, Leon Moodley<sup>3</sup>, Stig Westerlund<sup>3</sup>, Andrew Sweetman<sup>3</sup>, Dick Van Oevelen<sup>3</sup>, Katja Guilini<sup>1</sup>, and Ann Vanreusel<sup>1</sup>

<sup>1</sup> Marine Biology Research Group, Ghent University, Krijgslaan, Ghent, Belgium  
E-mail: Lisa.Mevenkamp@ugent.be

<sup>2</sup> Department of Ecosystem Studies, NIOZ Royal Netherlands Institute for Sea Research, Korringaweg 7, 4401 NT Yerseke, the Netherlands

<sup>3</sup> International Research Institute of Stavanger (IRIS), Randaberg, Norway

Mining activities in the deep-sea will have different impacts on marine benthic life on different scales including mechanical habitat destruction, release of toxic amounts of heavy metals or burial with sediment of mine tailings. We evaluated the short term effect burial with mine tailings on ecosystem functions of deep-sea (207 metres) benthos originating from a Norwegian fjord. Sediment cores were sampled and incubated in the laboratory under constant water flow and a temperature of 8°C. Mine tailings were added to the cores resulting in different thicknesses of added substrate (0cm (control), 0.1cm, 0.5cm and 3cm). Sediment cores were kept at constant temperature and water flow for a duration of eleven days. Oxygen profiles were measured at the beginning and at the end of the experiment. Sediment oxygen consumption, silicate content and dissolved inorganic carbon content was measured at the end of the incubation. Sediment cores were then sliced and macrofauna as well as meiofauna was sampled. In order to assess viability of the animals, a life check was done for macrofauna and meiofauna was stained with a vital stain, Trypan Blue. Since analysis of meiofauna is very time consuming, only preliminary results of the experiment will be given.